Arysta LifeScience Iberia celebrates the First Meeting of Fruit

Arysta LifeScience organized the first conference on Fruit, which took place on February 16 in the auditorium of the Agri-food Science and Technology Park of Lleida, in which the main products in the portfolio were presented in this crop.

The conference was opened by Josep Clotet, director of the Science Park, who thanked the realization of this event and invited the audience to better understand the facilities Parc. Then Clara Narrillos, Director of Marketing and Development Arysta LS, presented the new company and highlighted the major brands of the catalog.

Cruz Maria Puyal, territorial delegate in the Northeast area Arysta LS, presented the advantages of a good positioning in the treatment Syllit flow (dodina 40%) for the control of Venturia sp., Noting that Syllit flow has a unique mode of action and a great preventive and curative effect in controlling this disease.

- Preventive, because it destroys the target fungus spores before they can develop hyphae and penetrate the host plant.

- Healing, because it is a product able to enter into the parenchyma cells, and destroy the hyphae that are already within them.

Thus, the most appropriate treatment for the first time is swollen bud and must be repeated if high incidence of disease or in the case of an area where the disease is endemic.

Furthermore, during the presentation, demonstration images usability of the last incorporated in the portfolio of fruit, Malvin 80 WG product were projected (capture 80%), with an innovative formulation that exhibits high solubility and avoiding foaming in the tank.

Giancarlo Chiot, responsible European Development insecticides and acaricides for Arysta LS, said management Dimilin 25 WP (diflubenzuron 25%) for control Psylla spp. and Cydia pomonella. Dimilin 25 WP insecticide is a growth regulator that interferes with the synthesis of chitin present in the cuticle of insects, affecting the molting process. Controls, therefore the plague, acting on the egg and larval stages, interrupting their life cycle and avoiding the appearance of adults.

Chiot Giancarlo stressed that a good product positioning depending on the target pest is crucial, so that, for control of Cydia pomonella, treatment must take place immediately before the oviposition time, while for control Psylla spp the optimal efficiency is obtained with treatments performed after commissioning, with a presence of white or above the yellow eggs eggs. He also stressed the good toxicological profile Dimilin 25 WP, being a safe product for users, farming, poultry, fish and beneficial insects.

Xavier Miarnau, researcher at the Experimental Station of the IRTA of Lleida, presented the main results of the development project Syllit flow in the cultivation of almonds, as part of the strategy to control ocher stain.

At the end of the day, Arysta LS cocktail for all attendees in the same Science Park offered.

Source: Arysta LifeScience Iberia